

12-20-2020

CHARACTERISTICS OF THE FUNCTIONAL STATE OF THE PSYCHOEMOTIONAL STATE IN PATIENTS WITH CHRONIC OBSTRUCTIVE PULMONARY DISEASE AFTER CORONAVIRUS INFECTION AND THE EFFECTS OF COMPLEX THERAPY

Sh.Sh. Tilloeva

Republican Specialized Scientific and Practical Medical Center of Therapy and Rehabilitation Republic of Uzbekistan

D.A. Rakhimova

Bukhara State Medical Institute

Ch.Sh. Bokiyeva

Bukhara State Medical Institute

Follow this and additional works at: <https://uzjournals.edu.uz/pediatrics>

Recommended Citation

Tilloeva, Sh.Sh.; Rakhimova, D.A.; and Bokiyeva, Ch.Sh. (2020) "CHARACTERISTICS OF THE FUNCTIONAL STATE OF THE PSYCHOEMOTIONAL STATE IN PATIENTS WITH CHRONIC OBSTRUCTIVE PULMONARY DISEASE AFTER CORONAVIRUS INFECTION AND THE EFFECTS OF COMPLEX THERAPY," *Central Asian Journal of Pediatrics*: Vol. 2020 : Iss. 4 , Article 5.

Available at: <https://uzjournals.edu.uz/pediatrics/vol2020/iss4/5>

This Article is brought to you for free and open access by 2030 Uzbekistan Research Online. It has been accepted for inclusion in Central Asian Journal of Pediatrics by an authorized editor of 2030 Uzbekistan Research Online. For more information, please contact sh.erkinov@edu.uz.

**CHARACTERISTICS OF THE FUNCTIONAL STATE OF THE
PSYCHOEMOTIONAL STATE IN PATIENTS WITH CHRONIC
OBSTRUCTIVE PULMONARY DISEASE AFTER CORONAVIRUS
INFECTION AND THE EFFECTS OF COMPLEX THERAPY**

Tilloeva Sh.Sh., Rakhimova D.A., Bokiyeva Ch.Sh.

Republican Specialized Scientific and Practical Medical Center of Therapy and
Rehabilitation Republic of Uzbekistan
Bukhara State Medical Institute

Resume

To study the quality of life, the psychoemotional state and the effect of complex therapy on the parameters of the ventilation-perfusion function of the lung and the parameters of the diastolic function of the right ventricle of the heart in patients with chronic obstructive pulmonary disease after coronavirus infection.

Key words: chronic obstructive pulmonary disease, coronavirus infection, right ventricular hypertrophy, vascular endothelial function, diastolic dysfunction of the right ventricle

**ХАРАКТЕРИСТИКА ФУНКЦИОНАЛЬНОГО СОСТОЯНИЯ
ПСИХОЭМОЦИОНАЛЬНОГО СОСТОЯНИЯ У БОЛЬНЫХ ХРОНИЧЕСКОЙ
ОБСТРУКТИВНОЙ БОЛЕЗНИ ЛЕГКИХ ПОСЛЕ КОРОНАВИРУСНОЙ
ИНФЕКЦИИ И ЭФФЕКТЫ КОМПЛЕКСНОЙ ТЕРАПИИ**

Тиллоева Ш.Ш., Рахимова Д.А., Вокиева Ч.Ш

Республиканский специализированный научно-практический медицинский
центр терапии и медицинской реабилитации Республика Узбекистан

Бухарский государственный медицинский институт

Резюме

Изучить качество жизни, психоэмоциональное состояние и влияние комплексной терапии на показатели вентиляционно-перфузионной функции легкого и параметры диастолической функции правого желудочка сердца у пациентов с хронической обструктивной болезнью легких после коронавирусной инфекции.

Ключевые слова: хронической обструктивной болезнью легких, коронавирусная инфекция, гипертрофия правого желудочка, эндотелиальной функции сосудов, диастолической дисфункцией правого желудочка (ПЖ), амлодипин.

КОРОНАВИРУС ИНФЕКЦИЯСИДАН СЎНГ СУРУНКАЛИ ОБСТРУКТИВ ЎПКА КАСАЛЛИГИ БИЛАН ОҒРИГАН БЕМОРЛАРДА ПСИХОЭМОЦИОНАЛ ҲОЛАТНИНГ ФУНКЦИОНАЛ ҲОЛАТИ ХУСУСИЯТЛАРИ ВА КОМПЛЕКС ДАВОНИНГ ХУСУСИЯТЛАРИ

Тиллоева Ш.Ш., Рахимова Д.А., Вокиева Ч.Ш

Республиканский специализированный научно-практический медицинский
центр терапии и медицинской реабилитации Республика Узбекистан
Бухарский государственный медицинский институт

Резюме,

Тадқиқот мақсади ўпканинг сурункали обструктив касаллиги бўлган беморларда коронавирус инфекциясидан кейин ўпка гемодинамикаси, юрак ўнг бўлмача диастолик функцияси кўрсаткичларини комплекс терапия муолажалари таъсирида ижобий силжишни ўрганиш. Натижаларга кўра, ўпканинг сурункали обструктив касаллиги оғриган беморларда базисли даво билан биргаликда кальций канали блокаторлари билан биргаликда қўллаш ўпка гемодинамикасининг яхшиланишига олиб келади.

Калит сўзлар: ўпканинг сурункали обструктив касаллиги, коронавирус инфекцияси ўнг қоринча гипертрофияси, томирларнинг эндотелиал функцияси, ўнг қоринча диастолик дисфункцияси, амлодипин.

Relevance

Chronic obstructive pulmonary disease (COPD), as a cause of death, ranks 4th in the world in the age group over 45 years old and is the only disease in which mortality continues to increase. Due to the widespread prevalence of COPD, direct medical and indirect costs associated with morbidity and premature mortality pose a serious economic and social problem for society in general and health authorities in particular [1,2].

Obviously, only in-depth studying all aspects of this disease will help find a way out of the current impasse. Chronic obstructive pulmonary disease is one of the leading diseases leading to disability, and tends to a steady increase in primary cases in both industrialized and developing countries [3,8]. The risk of COPD varies from 1.85 to 2.88 per 1,000 population. According to the conclusion of the World Health Organization, in the last 10 years, COPD among the causes of death is in 4th place, being the cause of 4.9% of deaths of all diseases. According to experts' forecasts, by 2020, COPD will take 3rd place in the structure of total mortality [7,8,26]. The acute hypoxemic patients may experience dyspnoea that may persist despite the

administration of oxygen flows >10-15 L/min with a reservoir mask [3,15]. In these cases, other devices, such as High-Flow Nasal Oxygen (HFNO) or the application of Continuous Positive Airways Pressure (CPAP) or Non-Invasive Ventilation (NIV) may be useful. However, it is important to point out that these interventions have to be used only in appropriate hospital settings in order to be prepared for a more aggressive treatment. When indicated, the administration of CPAP/NIV can be carried out with various interfaces, depending on the availability and indications (oro-nasal mask, total face or helmet). When using CPAP/NIV, it is important to consider the potential environmental diffusion of aerosol particles of the virus [12,25]. In particular, one of the critical issues of the SARS-Cov-2 in the intermediate phase (between the onset of disease and potential critical evolution, also in relation to comorbidity) is represented by the selection of oxygen therapy and the level of care. Non-invasive supports (CPAP, NIV and HFNO) can correct hypoxemia and respiratory failure (even in the absence of clear evidence from proper randomized controlled studies), delaying or avoiding endotracheal intubation (and its potential complications and effects on the outcome) [14,26]. However, looking at SARS epidemic data, physiotherapists have to be careful when treating these patients because there is evidence that NIV may increase the risk of aerial spreading of the virus [11,16]. Thus, if a patient shows prognostic factors suggesting the need of invasive ventilation [17,27,16], it is preferable to carry out elective intubation, rather than emergency intubation in critical conditions.

Along with smoking, industrial pollutants, such as dust, gases, smoke, toxic aerosols, are becoming an important risk factor for the development of COPD, which significantly increase the frequency and severity of the disease [9]. The data given in the literature by domestic and foreign authors indicate that about 15–20% of all cases of COPD are associated with professional activity [5, 4,9].

Chronic obstructive pulmonary disease is essentially a disease with features of a destructive process, as a result of which damage to the entire respiratory system develops, including the bronchi up to the terminal bronchioles, parenchyma and interstitium [1,10]. An important feature of the pathological process in COPD is its low

reversibility [2]. The main method of treatment for such patients has long been the use of bronchodilators, starting with belladonna. However, as new knowledge about the essence of pathology was gained, new approaches to therapy were developed [8]. First of all, it became clear that COPD patients are not the same. They differ in the course of the disease, in symptomatology, in prognosis, in response to one or another therapy, which is defined as phenotypes [4,7,8].

The aim of the study was to study pulmonary hemodynamics and diastolic dysfunction of the right ventricle (RV) in patients with chronic obstructive pulmonary disease COPD in dynamics, combination therapy with calcium channel antagonists is recommended to improve pulmonary hemodynamics and bronchial patency[14,7,18].


Materials and methods

Thirty-five patients with COPD and 20 healthy individuals (AP) were examined. In patients, according to Doppler echocardiography parameters, PH (the level of mean pulmonary arterial pressure $LAPav > 25$ mm Hg) and the presence of RV without PH (thickness of the anterior wall of the RV according to EchoCG < 5 mm, with anteroposterior size of the RV > 2.5 cm). Patients were divided into 2 groups and a, b subgroups: 1a - gr. control group (CG) 8 patients with COPD with PH; 1b - column 7 patients with COPD with PDG received basic therapy (BT, GINA 2006), 2a - gr. 10 patients with COPD with PH; 2b - column 8 patients with COPD with PDG received BT and a combination of calcium antagonist dihydropyridine series amlodipine (A). For 10 days, patients received against the background of BT received a combination of a calcium antagonist of the dihydropyridine series amlodipine (An) 2.5-5 mg / day. FVC), forced expiratory volume in 1 sec (FEV1) and Tiffeneau's index (FEV1 / FVC); the study of peripheral blood flow-endothelium-dependent vasodilation (EDVD) was carried out with an ultrasound device SonoScape SSI-8000 (China); blood oxygen saturation (SaO₂) was assessed using the pulse oximetric method "OXY" (Germany); Doppler echocardiography was evaluated using a Vivid S60 ultrasound machine (Sweden) and SonoScape SSI-8000 (China) in accordance with the recommendations of the American Society. The results were processed using the Excel software package

using the Student's t-test. Differences between the studied parameters were considered significant at $p < 0.05$.

Results and discussion

Before treatment in conditions of increased myocardial oxygen demand in patients, there was a violation of the parameters of endothelium-dependent vasodilation: a decrease in the maximum blood flow velocity in the brachial artery after a reactive test (Vmax) - by 18, 9 (1st group) versus 28.4% (2nd); an increase in the index of circulatory resistance (ICR) - by 18.2 and 29.1% ($p < 0.001$ in relation to the control group). The data obtained correspond to the opinion of A.G. Chuchalin. [7] that the intensification of inflammatory phenomena leads not only to local changes, but also to significant systemic changes. Thus, disturbances in the peripheral mechanisms of regulation of vascular tone develop earlier and more significantly in patients with PH and COPD compared to COPD with PDG ($p < 0.05$, in relation to indicators of PL) (figure).



Picture. Parameters of EDVP and FVD in patients with COPD in the dynamics of complex treatment.

According to FVD, there was a decrease in the ventilation state of the bronchopulmonary system in all patients with PH and PDG. Thus, the FEV1 index in all patients was $-46.1 \pm 1.6\%$, ($p < 0.005$), SaO2, respectively, $89.6 \pm 1.4\%$ ($p < 0.05$), which is typical for an increase in bronchial obstruction. The results obtained revealed

the relationship between the clinical course, an increase in LAP avg. and ventilation and perfusion disorders in patients with COPD.

There was a tendency towards a more pronounced decrease in the atrial filling fraction (FPF,%) of the pancreas in patients with COPD 2 - gr. LH and RV of subgroups, respectively, by 1.05 and 1.08 times; and an increase in the time of deceleration of the maximum speed of early diastolic filling (VZ, m / s) by 1.07 and 1.15 times; a decrease in the ratio of early and late diastolic filling (E \ A) by 1.17 and 1.45 times, respectively ($p < 0.05$), which indicates an increase in myocardial stiffness and impaired passive relaxation.

After the therapy, there is an improvement in the studied parameters. Positive correlations were revealed between SaO₂: isovolumic relaxation time (VIR, m / s) and VZ, respectively, 0.40 and 0.35 ($p < 0.05$). After treatment in 2-gr. LH and RVD subgroups, in patients, the blood SaO₂ values increased to 89.8 ± 0.7 and $92.3 \pm 0.6\%$, respectively, by 1.10 and 1.16 times lower than the LP indicators ($p < 0.05$). In the dynamics of the complex treatment with Am E \ A, the indicators increased by 7.9 and 4.0%, respectively, and the mean pulmonary artery pressure (LAPav) by 1.20 and 1.16 times ($p < 0.01$). The use of BT alone does not have a significant effect on the parameters of pulmonary hemodynamics, providing only a decrease in LADav.

Correlation analysis showed that the increase in obstruction and hypoxemia are closely associated with the development of diastolic dysfunction of the pancreas. At the same time, the severity of the increase in LADav. has a significant dependence on the severity of COPD.

In our study, the use of amlodipine 2.5-5 mg / day against the background of BT in patients with COPD with PH led not only to a decrease in PAP avg., but was also accompanied by a positive shift in ventilation-perfusion lung capacity in group 2 of patients by 10.5% ($p < 0.01$). LAD Wed after treatment in group 2 decreased by 15.7% (significance of the difference with the indicators before treatment $p < 0.05$). It should be noted that in patients with COPD and PH with an increase in PAP avg. changes in the structure of filling the pancreas in diastole. The mutual aggravation of disorders in

the combination of cardiorespiratory pathology is based on the commonality of some links of pathogenesis - disorders of pulmonary and cardiac microcirculation, the development of hypoxemia and pulmonary hypertension. It was found that in patients with COPD with PH receiving amlodipine against the background of BT, there is an improvement in the parameters of pulmonary hemodynamics and ventilation capacity and bronchial patency of the lungs.

Conclusions

1. Violations of the peripheral mechanisms of regulation of vascular tone develop earlier and more significantly in severe clinical course of COPD with an increase in PAP cf. ($p < 0.05$).

2. Complex treatment of patients with COPD with pulmonary hypertension, including amlodipine, reduces the tone of pulmonary vessels and simultaneously improves endothelium-dependent vasodilation of peripheral vessels.

LIST OF REFERENCES:

1. Alyavi A.L., Rakhimova D.A., Sabirzhanova Z.T. Pulmonary hypertension / Monograph. - Tashkent, 2016.89C.
2. Akhmineeva A.Kh., Polunina O.S. Endothelial dysfunction in chronic obstructive pulmonary disease and bronchial asthma // Astrakhan Medical Journal. - 2012. - T. 7. - No. 3. - P. 43-46
3. Alyavi A.L., Rakhimova D.A., Sabirzhanova Z.T. Pulmonary hypertension / Monograph. - Tashkent, 2016.89C.
4. Akhmineeva A.Kh., Polunina O.S. Endothelial dysfunction in chronic obstructive pulmonary disease and bronchial asthma // Astrakhan Medical Journal. - 2012. - T. 7. - No. 3. - P. 43-46
5. Pulmonology. National leadership / ed. A.G. Chuchalin. -M.: "GEOTAR-Media", 2009. P.518.
6. Ubaidullaev A.M., Alyavi A.L., Rakhimova D.A. Parameters of psychovegetative regulation of heart activity in patients with chronic obstructive pulmonary disease

complicated by cor pulmonale // Therapeutic Bulletin of Uzbekistan, materials of the conf. - Tashkent, 2014. - No. 4. - P. 148.

7. Chuchalin A.G. and other Federal clinical guidelines for the diagnosis and treatment of chronic obstructive pulmonary disease. M., 2013
8. Global Strategy for the Diagnosis, Management and Prevention of COPD. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Updated 2014. <http://www.goldcopd.org>
9. Koblizek V. et al .; Czech Pneumological and Phthisiological Society // Biomed. Pap. Med. Fac. Univ. Palacky Olomouc Czech. Repub. 2013. V. 157. No. 2. P. 189.
10. Dretzke J. et al. // Health Technol. Assess. 2015. V. 19.No. 81. P. 1.
11. 11.Lai CC, Shih TP, Ko WC, et al. Severe acute respiratory syndrome
12. Coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int J Antimicrob Agents* 2020;55:105924. doi: 10.1016/j.ijantimicag. 2020.105924.
13. Wujtewicz M, Dylczyk-Sommer A, Aszkiełowicz A, et al. COVID-19 - what should anaesthesiologists and intensivists know about it? *Anaesthesiol Intensive Ther* 2020. pii: 40133. doi: 10.5114/ait.2020.93756.
14. Kallet R.H., Hemphill J.C., Dicker R.A., et al. The spontaneous breathing pattern and work of breathing of patients with acute respiratory distress syndrome and acute lung injury. *Respir Care* 2007; 52:989-95.
15. Meng L., Qiu H., Wan L., et al. Intubation and ventilation amid the COVID-19 outbreak: Wuhan's Experience. *Anesthesiology* 2020. doi: 10.1097/ALN.0000000000003296. [Epub ahead of print]
16. Seto W.H., Tsang D., Yung R.W., et al. Effectiveness of precau- [Monaldi *Archives for Chest Disease* 2020; 90:1285, 167 168 page.
17. Ding L., Wang L., Ma W., He H. Efficacy and safety of early prone positioning combined with HFNC or NIV in moderate to severe ARDS: a multi-center prospective cohort study. *Crit Care* 2020;24:28. doi: 10.1186/s13054-020-2738-5

18. Zhou F., Yu T., Du R., et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020. pii: S0140-6736(20)30566-3. doi: 10.1016/S0140-6736(20)30566-3.
19. Hui D.S., Chow B.K., Chu L., et al. Exhaled air dispersion and removal is influenced by isolation room size and ventilation settings during oxygen delivery via nasal cannula. *Respirology* 2011;16:1005-13. doi: 10.1111/j.1440-1843.2011.01995.x
20. Hui D.S., Chow B.K., Chu L.CY, et al. Exhaled air and aerosolized droplet dispersion during application of a jet nebulizer. *Chest* 2009; 135:648-54. doi: 10.1378/chest.08-1998
21. Writing Group for the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial (ART) Investigators, Cavalcanti A.B., Suzumura E.A., et al. Effect of Lung Recruitment and Titrated Positive End-Expiratory Pressure (PEEP) vs low PEEP on mortality in patients with acute respiratory distress syndrome: A randomized clinical trial. *JAMA* 2017; 318:1335-45. doi: 0.1001/jama.2017.14171
22. Favretto D.O., Silveira R.C., Canini S.R., et al. Endotracheal suction in intubated critically ill adult patients undergoing mechanical ventilation: a systematic review. *Rev Lat Am Enfermagem* 2012; 20:997-1007.
23. Guerin C., Reignier J., Richard J.C., et al. Prone positioning in severe acute respiratory distress syndrome. *NEJM*. 2013;368:2159-68.
24. Blackwood B., Alderdice F., Burns K.E., et al. Protocolized versus nonprotocolized weaning for reducing the duration of mechanical ventilation in critically ill adult patients. *Cochrane Database Syst Rev* 2010; CD006904.
25. Pena-Lopez Y., Ramirez-Estrada S, Eshwara V.K., Rello J. Limiting ventilator-associated complications in ICU intubated subjects: strategies to prevent ventilator-associated events and improve outcomes. *Expert Rev Respir Med* 2018; 12:1037-50.

26. Fan E. Critical illness neuromyopathy and the role of physical therapy and rehabilitation in critically ill patients. *Respir Care* 2012; 57:933-44; discussion 944-946.
27. Van Aerde N., Meersseman P., Debaveye Y., et al. Five-year impact of ICU-acquired neuromuscular complications: a prospective, observational study. *Intensive Care Med* 2020 doi: 10.1007/s00134-020-05927-5. [Epub ahead of print]
28. Ambrosino N., Makhabah D.N. Comprehensive physiotherapy management in ARDS. *Minerva Anestesiol* 2013; 79:554-63.

Entered 09.04.2021